

I. OVERVIEW

The Northeast Kingdom is recognized for its diverse wildlife, large undeveloped areas, and vast woodlands. The region's natural resources (depicted in Map 11 on the following page) provide residents and others a variety of benefits. The largest source of revenue in the region is from the harvesting, processing, and manufacturing of forest products; a wide variety of recreational activities are available in the forests and on lakes; and much of the tourism industry relies on the healthy and scenic environment to remain viable. Therefore, the natural resources in the Northeast Kingdom have intrinsic scenic and economic values that require careful consideration when making planning decisions. The overarching goal for the region is to balance local economic needs with the protection of the resources that so many of region's residents enjoy and depend upon.

The Northeast Kingdom lies mostly within three physiographic regions:


- *The Northeast Highlands*, an extension of New Hampshire's White Mountains, make up most of Essex County and northern Caledonia County. On average, this area is cooler than the rest of the state. The growing season here averages less than 90 days and snowfall accumulation frequently exceeds 36 inches.
- In much of Orleans County and parts of Caledonian County the topography is primarily *rolling hills* interspersed with occasional plains of fertile agricultural soils. Both of these physiographic regions have extensive glacial deposits.
- The third region is the *Connecticut River Valley*, which extends the length of the region along its eastern border. Level topography and rich alluvial soils well suited for agriculture characterize this physiographic region.

The forests are mainly northern hardwoods with large stands of red spruce and balsam fir. Black spruce and succession species such as white pine and aspen fill recent clearings. The region contains some of the State's largest bog and wetlands complexes, with fabulous stands of red pine, black spruce, hemlock and hardwoods dispersed throughout. Essex County has more wetlands than any other county in Vermont.

The majority of the region's water drains either north to Quebec as part of the St. Francois River watershed or east and south as part of the Connecticut River watershed. Much of the region's western edge drains north and west as part of the vast Lake Champlain basin. The region's lakes and rivers are famous for the excellent and diverse fishing opportunities they offer. The more than 130 lakes and ponds found concentrated in the region represent a disproportionately high share of the State's total. This region is home to most of Vermont's larger, deeper lakes and the legendary 20-30 pound trout that have inhabited them since the last ice age.

This combination of forest and water resources creates prime habitat for many wildlife species, and draws many to the Northeast Kingdom.

MAP 11:



II. WATER RESOURCES

Water Quality

According to the Water Quality Division of the Department of Environmental Conservation, the water quality of all rivers, streams, lakes, and ponds in Vermont is considered good. When mercury contamination in fish is added to the equation, the rating is fair. No comprehensive studies have been completed on the quality of Vermont's groundwater, yet based on the small number of public water supplies throughout the state that have detected any contamination, the Water Quality Department describes Vermont's groundwater as "excellent."

Vermont's classification system for surface water establishes management goals and practical uses. The Water Resources Board classifies all surface waters in Vermont as either Class A or Class B. The July 2, 2000 Water Quality Standards recognize two categories of Class A waters. Class A(1) waters are ecological waters, which are managed to maintain waters in a natural condition. All of the waters above 2,500 feet in elevation are classified A(1). Class A(2) waters are managed as public water supplies and therefore allow moderate water level fluctuation. Class B waters are designated as being either Water Management Type 1, 2, or 3 depending upon their protection and management. An overlay of both Class A and Class B waters is an Outstanding Resource Water (ORW). These waters are designated by the Water Resources Board as having exceptional natural, recreational, cultural, or scenic value. Most waters in the Northeast Kingdom, like in the rest of Vermont, are Class B, which is consistent with State policy to achieve and maintain Class B waters with suitability for swimming, boating, and drinking with treatment as well as for irrigation and livestock watering.

The Water Quality Division maintains a river and lake assessment database listing impaired surface waters. This database is updated every two years with the most recent information and data from a variety of sources. Table 9.0 displays the impaired lakes and rivers within the Northeast Kingdom from this database. Threats listed are addressed at the local, regional, state or federal levels. For instance, agricultural runoff is a local and regional issue, whereas sewage overflows and mercury issues should be addressed across the entire range of management levels.

| Table 9.0: State Targeted-Impaired Surface Waters. | |
|--|---|
| Water Body | Threat or Impairment |
| Levi Pond – Groton | Extremely sensitive to acidification; episodic |
| Stearns Brook & Tributary – Holland | Undefined: Agricultural runoff |
| Lake Salem – Derby | Elevated levels of mercury in Walleye |
| Crystal Brook – Derby | Sediment & nutrients: Agricultural runoff |
| Ticklenaked Pond – Ryegate | Phosphorus: Algae blooms, high pH, low D.O. |
| Lake Memphremagog – Newport | Phosphorus: Excessive algae growth, nutrient enrichment |
| South Bay – Newport | Phosphorus: Nutrient enrichment, nuisance algal |
| Comerford Reservoir – Barnet | Elevated levels of mercury in all fish |
| Moore Reservoir – Waterford | Elevated levels of mercury in all fish |
| Lower Sleepers River – St. Johnsbury | E. Coli: St. Johnsbury WWTF Collection System passes combined sewer |
| Passumpsic River from Pierce Mills to below Passumpsic Dam | E. Coli: St. Johnsbury WWTF Collection System passes combined sewer overflows |

Source: VT DEC-WQD,303(d) List of Impaired Surface Waters in need of TMDL, 2004

Watersheds

A watershed is a geographic area in which all water flows into a single river. Boundaries of watersheds are therefore the highest points of elevation surrounding the water body. Watersheds can be small and localized, or large enough to incorporate many drainage areas flowing into a vast water body. Vermont is divided into 17 major watersheds. The three counties of our region contain waters that flow within the St. Francois or Lake Memphremagog; Passumpsic; Upper Connecticut; Lamoille; Missisquoi; Stevens, Wells, Waits, Ompompanoosuc or "Basin 14"; and Winooski watersheds.



The Agency of Natural Resources should complete a watershed basin planning process for all 17 major watersheds by the year 2006. Each basin will have a watershed council comprised of area residents and technical advisors to create a comprehensive watershed plan. Plans should contain present water quality conditions, problems of local concern, management goals and implementation strategies. The Lamoille River Watershed Council has already been established and is expected to complete its plan by 2005. The Basin 14 Council has been established which includes the subgroups of the Stevens River Watershed Council and the Wells River Watershed Council. This plan is not expected to be complete until 2006.

Joint working groups were formed to address agricultural issues and lake/river water quality monitoring and objectives. The Lake Memphremagog Quebec/Vermont Working Group, established by the governments of Quebec and Vermont in 1989, completed their *Final Report on Managing Lake Memphremagog and its Environment* in 1993. Recommendations were made to reduce point and non-point sources of pollution, and a Vermont Memphremagog Steering Committee was formed in the spring of 1995 to work with its Quebec counterpart to implement priority recommendations.

The Connecticut River Joint Commissions were established by the legislatures of New Hampshire and Vermont and directed to serve in an advisory capacity, promoting public involvement in decisions affecting the Connecticut River and its watershed. Five local river subcommittees operate along the river in Vermont and New Hampshire. The two subcommittees in our region are the Headwaters and the Riverbend. The Connecticut River Watershed Council is another citizen-based advocate for the environmental well-being of the Connecticut River. This Council, founded in 1952, includes representatives from the four states where the river flows: Vermont, New Hampshire, Massachusetts and Connecticut. This organization conducts research, produces publications, awards small project grants, and initiates programs to generate local activism in the River watershed.

There are also several local Lake Associations involved in watershed activities throughout our region. One example is the Lake Parker Association in West Glover, which has done work with watershed assessment, road erosion control, shoreland vegetation enhancement, lay monitoring and invasive aquatic species prevention. These lake associations have created a network for local citizens to help restore the health of their water systems.

MAP 12:



NVDA Region: Watersheds

Table 9.1: Local Lake Associations

| Local Associations involved in Watershed Activities: | Programs Conducted by Local Lake Associations: |
|--|---|
| Lake Memphremagog Watershed | |
| Averill Lakes Assoc. (Great and Little Averill, Forest Lake) Crystal Lake Reservation Assoc. Daniel’s Pond Assoc. Echo Lake Assoc. Elligo Lake Assoc. Friends of Little Hosmer Island Pond Assoc. Lake Parker Assoc. Memphremagog Conservation, Inc. Salem Lake Assoc. Seymour Lake Assoc. Shadow Lake Assoc. Westmore Assoc. (Lake Willoughby, Long Pond, Bald Hill Pond) | ⇒ Bacteria Monitoring ⇒ Exotic Species Spread Prevention ⇒ Lake Assessment ⇒ Lake Protection Through Town Zoning ⇒ Land Conservation ⇒ Landowner Education ⇒ Lay Monitoring Program ⇒ Local Advocacy ⇒ Road Erosion Control ⇒ Runoff Diversion ⇒ Shoreland Vegetation Enhancement ⇒ Testing of Shoreline Septic Systems ⇒ Watershed Assessment |
| Lamoille River Watershed | |
| Greensboro Assoc. (Caspian Lake) | ⇒ Watershed Management Committee |
| Passumpsic River Watershed | |
| Cole’s Pond Assoc. Joe’s Pond Assoc. Lyford Pond Assoc. South End Newark Pond Assoc. | Basin 14: SEWER Save Everyone’s Wells River Passumpsic River Network Passumpsic Valley Land Trust |
| Stevens River Watershed | CT River Watershed Advisory Commission |
| Harvey’s Lake Assoc. | Barton River CPP Program |
| Upper Connecticut River Watershed | Lamoille River Anglers Association |
| Maidstone Lake Assoc. Miles Pond Assoc. Neal Pond Assoc. | Missisquoi River Basin Association |

(VT Agency of Natural Resources, Water Quality Division, Pamphlet: “Watershed and Lake Associations of Vermont”, October 2003)

Surface Waters

The Northeast Kingdom has the largest amount of surface water of any region in Vermont, excluding Lake Champlain. The region is famous for its pristine trout and salmon lakes. Lakes such Willoughby, Seymour, and Caspian offer high quality trout fishing, while Lake Memphremagog and the Moore Reservoir are known for small-mouth bass. The Clyde, Black, Barton and Willoughby Rivers, primary tributaries to Lake Memphremagog, draw anglers from all over the Northeast for the large leaping rainbow trout and landlocked Atlantic salmon that migrate upstream from the big lake. Efforts to restore Atlantic salmon are taking place in the upper Connecticut River basin waters, such as the Passumpsic, Moose, Stevens and Wells Rivers. The region’s lakes and rivers serve as significant sources of recreation for residents and visitors, as well as being the source of drinking water for some communities.

Ground Water

Ground water is a critical resource for the rural areas of Vermont. Approximately 60 percent of Vermont's citizens depend upon ground water for drinking and general uses. According to the state Water Quality Division, in many rural communities nearly 100 percent of the public and private drinking water sources are from ground water. Ground water occurs in two general hydrogeologic settings, bedrock and unconsolidated aquifers. Igneous and metamorphic crystalline bedrock along with carbonate bedrock form the bedrock aquifers within the state. Unconsolidated deposits are comprised of glacial till, which is basically sand and gravel.

As ground water moves through these materials, the organic and mineral substances that are dissolved or picked up dictate the quality of water. These water sources often tend to be better quality than surface water because of the leaching process. However, toxic substances can stay with ground water for very long distances. Although groundwater quality is generally good, the resource is nonetheless fragile. Contaminated wells destroy property value. The cost of developing and installing new groundwater sources for public water supply is estimated between \$500,000 and \$1,000,000 (Water Quality Division, 2000). This cost is prohibitive for many towns in the Northeast Kingdom, making prevention and education critical management tools. One way to avoid costly groundwater contamination cleanup is to begin a Well Head Protection Program (WHPP). This is addressed in the Water Supply section of the Utilities and Facilities chapter of this document.

Wetlands

Vermont's wetlands are defined as those areas of the state that are inundated by surface or ground water with a frequency sufficient to support plants and animals that depend on saturated or seasonally saturated soil conditions for growth and reproduction. These areas are commonly known as ponds, bogs, fens, marshes, wet meadows, shrub swamps, and wooded swamps. Wetlands often occur in association with lakes, ponds, rivers, and streams, creating transitional areas between dry land and open water. However, wetlands can also be isolated from any obvious connection to surface water. In order to be classified as a wetland under Vermont law, an area must have wetland soils and wetland plants, in addition to at least seasonal water. Wetland soils are often anaerobic and the plants have adapted to growing in such waterlogged conditions.

Every town in the region contains wetlands that have been designated by the state as significant (see Table 4). The Vermont Wetland Rules classify all wetlands into one of three classes. Classes One and Two are considered "significant" and protected under the Vermont Wetland Rules. All three wetland types are protected by Vermont's Act 250.

Wetlands provide important ecological functions, including flood and erosion control, and providing habitat for fish and wildlife. They aid in the maintenance of water supplies by trapping nutrients and sediments and recharging groundwater, and they provide recreational and educational opportunities.

Wetlands were once considered wastelands and were thought to be “improved” by draining and filling. As a result, nearly half of Vermont’s original wetland resources have been lost, and hundreds of acres are lost annually. Human activities and development continue to encroach upon this finite resource. Agriculture and forestry activities as well as residential, commercial and industrial development all result in wetland alteration. Replacing new wetlands is costly and often impractical, so wetlands preservation is important.

Floodplains

Floodplain maps identify flood hazard areas, defined as those areas that are inundated by the waters of the 100-year flood. Limiting extensive development in flood-prone areas is the best way to prevent flood damage while allowing the floodplain to function as it should. Overdevelopment in flood hazard areas increases risks to human life, property damage, and leads to habitat loss for wildlife species. Preventative measures, such as flood hazard regulations, can reduce these risks. Flood hazard regulations enable property owners to obtain both flood insurance and mortgage loans for property in flood hazard areas. Overdevelopment in floodplains can lead to an increase in intensity and frequency of flooding. This is especially true in areas that are losing wetlands due to development. Wetlands act like sponges-soaking up excess water-and when these are filled in or altered, there is an increased potential for flooding along other parts of the stream banks.

Prevention is the best strategy for reducing environmental and human impacts from flooding. Preventative strategies include various planning and zoning tools, open space preservation, and watershed management.

Pollution

The Connecticut River Joint Commissions’ *The Watershed Guide* indicates that most aquifer contamination comes from “non-point” sources. This means that contaminants can come from drainage areas some distance away from the point of contamination. Contamination sources include “point” sources such as leaking gasoline storage tanks, failing septic systems, salt storage piles, landfills, storage tanks, or “non-point” sources such as urban and agricultural runoff, and mining operations. Cleanup of bedrock aquifers is costly and often ineffective.

Many farmers use Best Management Practices (BMPs) to reduce the amount of pollution entering waterways. Millions of dollars in state funds have been committed to help farms implement Best Management Practices. These practices are also useful for landowners and towns to avoid polluting waterways. There are BMPs for timber harvesting, construction and development, septic system care, road construction, road salting and dumping, golf courses, site excavation, sand and gravel operations, urban runoff, chemical and petroleum products, land application of bio-solids, and docks, moorings, and marinas. Education and voluntary compliance are essential to mitigating the production of non-point source pollution.

Erosion and Sedimentation

Erosion and discharge of sediment into rivers, lakes, and streams can cause significant damage to aquatic communities. Sediments suspended in the water or deposited on bottoms can

adversely affect the growth, feeding, and reproduction of many organisms. Other impacts include loss of topsoil, contamination of water by heavy metals, increased flooding potentials and high municipal costs for ditch, culvert, and drain cleanup. The rate of erosion is influenced by the type of soil, vegetative cover, topography, and climate.

Vermont's extensive networks of dirt roads are a major cause of erosion and sedimentation when improperly built and maintained. It is important that the state, towns, and landowners cooperate in efforts to maintain these roads with better practices to protect water quality and save funds over the long term. The Better Backroads Program, established by the Agency of Natural Resources, is a popular program for education on appropriate road maintenance.

Failing Septic Systems

Failing septic systems can be a major source of pollution. Septic systems fail due to inadequate soils, poor design or construction, inadequate maintenance, or increased use from seasonal to year-round use. Failing septic systems can result in either effluent surfacing on the ground or contamination of ground water. Both situations contaminate water supplies and are hazardous to human health.

Water Quality

As development around the region's lakes has increased, so has the use of these lakes and the amount of pollution entering the water. A lake can naturally treat a certain amount of pollution but if the amount surpasses the lake's natural ability to treat the waste, it starts to accumulate in the sediments and water column. The Northeast Kingdom's lakes are cold water lakes which have a much harder time breaking down pollutants. Once a particular concentration of nutrient pollution is reached, there may be a decrease in the amount of oxygen available for fish along with an increase in lake temperatures. Sediment build-up and algae growth are natural processes in the life cycle of a lake. However, when phosphorous and sediments enter a lake through erosion, agricultural run-off, or failed septic systems, this process is accelerated dramatically, hastening the lake's decline.

A decline in water quality may affect the real estate value of lakefront properties. Vermont has no statewide shoreline zoning laws, and much of the shoreline development currently taking place receives little or no review. Water quality protection is dependent on municipalities enacting and enforcing good resource protection measures.

III. MINERAL RESOURCES

Soils

In general, soils can be classified as clay, silt or sand. Combinations of one or more of these can create many variations of soil. Soil is influenced by the organic matter that is deposited on the surface and by the organisms that exists within it, in combination with parent materials. Within soils, organisms and fungi provide food for animals and create organic matter for more efficient vegetative production. This vegetative layer, in turn, helps to purify surface water.

The availability of soils suitable for cultivation plays a crucial role in agricultural productivity. Out of the 257,000 acres of potentially primary agricultural soils in the Northeast Kingdom, 133,565 acres were in production in 1992 as compared to 142,832 acres in 1987. It should be noted that less than 8% of these acres are found in Essex County, where most of the terrain and soil composition is better suited for the forestry industry. Despite this fact, there are still important agricultural soils in Essex County along the Connecticut River.

Sand and Gravel

The primary sources of sand and gravel follow streams and waterways or are adjacent to water bodies. During the ice age streams tunneled through ice sheets. These streams allowed deposits of sand and gravel and other debris to build up, creating deposits called eskers. Water flow is easier through these deposits, forming rivers and streams. The Passumpsic River in Caledonia County is one of the largest, continuous eskers in the region, approximately 24 miles in length. Clear, clay-free materials of eskers are excellent for concrete and asphalt aggregates, roadbeds and other construction uses.

Sand and gravel deposition also happened when large glacial ice remnants melted. Much of the remaining sand and gravel was deposited in the valleys along waterways. One of the largest areas of sand and gravel deposits in the region extends from west of Island Pond to the eastern border of the Nulhegan Basin.

Sand and gravel deposits often serve as important areas for groundwater aquifer recharge and filtration. Disturbance of these areas can result in a reduction of their natural ability to retain and filter groundwater. As minerals are extracted, deposits become shallower and less able to filter contaminants from the water. Because of these infiltration concerns, the distances from gravel pits to surface and ground water supplies should be examined.

The prospects for sand and gravel extraction are difficult to measure. While the 1990 Census identified only three sources of sand and gravel extraction activity, the District 7 Act 250 Office has issued over 50 permits in the Northeast Kingdom counties for extraction of sand and gravel. In addition to these, there are small pits like many town sources for local road maintenance not subject to Act 250 standards. The contribution of mineral extraction to the local economy is difficult to ascertain because most extraction is done as part of some other business or is additional income for entrepreneurs. There are active granite quarries in Sheffield, Ryegate and South Ryegate.

Soil Compaction

Compacted soil, which occurs naturally, as well as through land development and industrial processes, makes it more difficult for water to be absorbed. This creates two changes to the soil formation process. First, water cannot flow through the soils in order to leach contaminated particles. Second, it creates erosion and carries away soil. As discussed under water resources, erosion contributes to flooding, removal of productive topsoil, distribution of chemicals on the soil, and sedimentation of surface water.

IV. AIR QUALITY

The air we breathe is less tangible than other resources, but equally important. Air quality is adversely affected by industrial, residential, and transportation emissions. The cyclical patterns of air are intricately connected with all other biological systems where change in one affects the others. Although the Northeast Kingdom is the antithesis of cityscapes where industrial pollution may be visible on a daily basis, there are still sources of air pollution that persist.

Industrial Emissions

Toxins such as sulfur dioxide and nitrogen oxide, emitted into the atmosphere when fossil fuels are burned, contribute to the acidification of our surface waters. The sulfates and nitrates remain in the atmosphere until rain transports them to the earth's surface where they increase the acidity of the soil and water. A study conducted for the Vermont Department of Environmental Conservation's Air Pollution Control Division concluded that 99.9 percent of the pollutants responsible for wet sulfate deposition in Vermont originate from out-of-state industrial sources.

Residential Emissions

The number one generator of air pollution in Vermont is the automobile, and its use continues to grow. Issues related to automobiles are addressed in the Transportation section of this regional plan. Household emissions include heating systems using wood and fossil fuel, and trash burning. Although municipalities no longer burn trash, there are many households that continue to burn residential wastes. The dioxins, hydrochlorides, carbon monoxide and various carcinogens released into the atmosphere pose risks to our health, the health of the surrounding ecosystems and create obnoxious odors for numerous neighbors.

Light Pollution

As development increases, outdoor lighting may become an issue for some residents. Neighbors of new developments with significant outdoor lighting can be particularly sensitive this. Directing outdoor lighting in new developments to reduce excessive light reduces complaints and allows people to enjoy the night sky.

V. WILDLIFE HABITATS

All wildlife species have three basic needs for survival: food, water and cover. The spatial relationship of these factors and their availability comprise the habitat of a given species. To promote a diversity of wildlife species, it is important to conserve various habitat types as well as critical areas that support basic needs for some species. For example, white-tailed deer live in a variety of forested and non-forested areas, but specific softwood wintering areas are critical for their survival. The deer have adapted to this habitat for their survival and without it they would not survive the harsh winters in Vermont.

Almost every kind of human development results in the loss of some wildlife habitat. Single developments, even at a large-scale, do not usually destroy a wildlife population. It is the cumulative impact of developments that gradually diminish wildlife habitats. It is difficult to plan for the

protection of wildlife habitat because their ranges tend to cross human defined political boundaries. This may require municipalities to work together on habitat protection issues.

The region's large tracts of undeveloped land provide excellent habitat for a variety of species. Both residents and visitors enjoy the variety of wildlife present. A 1996 survey by the Vermont Department of Fish and Wildlife revealed that 242,000 Vermont residents 16 years and older engaged in fishing, hunting, or wildlife-watching activities. In the same year, both resident and nonresidents spent \$341 million on wildlife-associated recreation in Vermont. This is a significant portion of our economic base and warrants attention.

Deer Wintering Areas

One of the most prized game species; the Northern White-tailed deer is very adaptable and thrives in this area due to the diversity of landscapes. The home range of white-tailed deer includes many habitats such as edges between fields and forests, wetlands, and broadleaf and needle-leaf forests. During the winter, it is critical for deer to stay in forested areas sheltered by needle-leaf softwoods such as balsam fir, cedar, spruce and white pine. These evergreens intercept the snow and create a refuge for the deer. Such a canopy offers thermal protection and greater mobility on the ground in deep winter. Although these "wintering areas" may only be a fraction (10%) of their yearly range, they are the single most important factor in determining the carrying capacity of the land. Without such habitat, the deer population would virtually die off.

Black Bear Habitat

Unlike deer wintering habitat, the Department of Fish and Wildlife has not completely mapped the critical black bear habitat in Vermont. Bear require large uninterrupted tracts of forestland that often contain American beech stands, wetlands and high elevations. In other areas, black bear populations have dwindled due to habitat loss resulting from highway and urban development. Unlike the wild turkey, black bear in Vermont were able to escape to high elevations during the 1800s when forested land was scarce. Habitat changes returned the black bear population, some say, to where it was before European settlers arrived. The black bear population in Vermont is now estimated at 3000. Public support, increased awareness of construction impacts on habitat, and private responsibility are necessary to protect the remote and extensive forestland necessary for the bear's survival.

River and Riparian Habitat

Vermont's river corridors provide critical habitat for endangered or threatened plants and animals. Along the rivers of the state, there are 27 species included on the Federal Endangered Species List. River and riparian habitats serve important functions for many plants and animals.

A riparian buffer is a unique ecosystem. It provides food and shelter for everything from caddis flies to brook trout to food for mammals and birds such as the river otter and kingfisher. The mink, bald eagle, Louisiana water thrush, dusky salamander, black bear and wood turtle all require streamside woods. These areas provide vital food supplies and safe

corridors for game species as well. Much of the river's natural riparian forest in Vermont is gone, removed for farming, development, or landowner access to the river.

Maintaining and repairing riparian buffers is inexpensive and can provide many economic benefits. A municipality will spend more money on bank stabilization, stormwater control and water quality improvements than it would by leaving or replanting riparian vegetation. Riparian buffers, which can eliminate the need for costly riprap, are an effective tool to avoid the costs of bridge collapse, and washed out roads. Farmers sometimes lose land when they clear riparian vegetation to grow crops because unstable banks can create a situation during floods when the river jumps its channel and cuts a new one into the cropland.

Travel Corridors and Contiguous Forestland

Many species, such as bobcat and black bear, use different areas dependent upon the season and they need ways to travel between habitats. Wildlife corridors enable their movement and hence their survival. Black bear require contiguous remote forestlands to survive. Black bear and bobcat are also considered "umbrella" species. This means that if migration corridors are effectively maintained to support these species, many other species with smaller home ranges will also benefit.

The division of habitat into smaller and smaller areas due to human settlement is termed "fragmentation." The transformation results in the direct loss of important habitat by the creation of gaps or barriers between isolated habitat blocks, rendering the habitat unusable for some animal species. The smaller the habitat block, the fewer the number of species can survive there and the less species diversity. Many species of wildlife rely on natural connections (corridors) between habitats to allow safe access within their habitat range.

Threatened and Endangered Species

The Endangered Species Listing by the U.S. Fish and Wildlife Service shows that there are six animal species and two plant species in our region whose survival is in question. Threatened species include the Bald Eagle, Canada Lynx, and Puritan Tiger Beetle. Endangered species include the Indiana Bat, Eastern Puma, Dwarf Wedge Mussel, Jesup's Milk-Vetch and the Northeastern Bulrush.

Often the preservation of these threatened and endangered species has more to do with habitat protection than any other factor. A species is considered endangered if it normally occurs in the state, and its continued existence appears to be in jeopardy. A species is threatened if its numbers are significantly declining due to loss of habitat or human disturbance. Habitat loss and degradation is the principle cause of the decline in biological diversity and is the number one factor relating to species endangerment. Human development is the primary cause of habitat loss and degradation.

Non-native Invasive Species

Many non-native species introduced to this region intentionally or otherwise are creating significant problems in both terrestrial and aquatic ecosystems. Because they have been

imported here from outside the region, there are no natural predators to control the spread of these non-native species. They often take over an area by out-competing the existing flora or fauna, choking out native species.

Purple Loosestrife and Eurasian Water milfoil are two invasive aquatic plants that currently infest a number of lakes and wetlands in our region. Purple Loosestrife is a wetland perennial that has infested thousands of acres in Vermont and can be found in almost every town in our region. It chokes out the native vegetation and can decrease the food and shelter for native wildlife species. Eurasian Water milfoil has also taken hold in several lakes of our region, including Brownington and Clyde Ponds and Lakes Elligo, Salem, Crystal, Willoughby, and Memphremagog. This freshwater seaweed creates a dense, impenetrable canopy of stems and leaves that consumes large quantities of oxygen, suffocating fish and other creatures. Preventing its expansion to other water bodies through education and controlling it within each lake's boundaries is currently the most effective way to manage it.


Open Space

The Northeast Kingdom is composed of rolling hills, farmlands, lakes and rivers, forests, country roads, and compact village centers. These areas combined create an open, picturesque landscape unlike any other. Open space provides not only scenic beauty and wildlife habitat, but is necessary for the numerous outdoor activities enjoyed by the region's residents and visitors, and is key to the agricultural and forestry traditions of the region. The region contains more than 1,300,000 acres of land. Almost 200,000 acres are either publicly owned or have public recreation/access easements. Many recreational activities rely on private landowners allowing access to their properties, so it is the responsibility of users to respect the landowner and their land. Vermont landowner liability law (12 V.S.A. 5793) maintains "an owner shall not be liable for property damage or personal injury sustained by a person who, without consideration, enters or goes upon the owner's land for a recreational use unless the damage or injury is the result of the willful or wanton misconduct of the owner." Still, according to the Vermont Department of Forests, Parks & Recreation, posting of private land in the state doubled in the last decade from approximately 100,000 acres in 1988 to approximately 250,000 acres in 1997.

Public Lands

The region contains many conserved public lands. Recently, more than 132,000 acres of remote forestland, primarily in Essex County, was conserved by Vermont's largest land conservation project. Of this, 84,000 acres was resold to Essex Timber Co. LLC, with easements to ensure that these lands are conserved as a working forest for the sustainable production of wood products as well as to maintain public access. In the same transaction, U.S. Fish and Wildlife Service formed the Silvio O. Conte National Wildlife Refuge in the towns of Lewis, Ferdinand, Bloomfield and Brunswick totaling nearly 28,000 acres. The 23,000 acre West Mountain Wildlife Management Area was created in this land transfer, as well. The goals of this purchase were to protect public access to the land; conserve and protect biological diversity, wildlife habitat and natural communities; and conduct sustainable management and utilization of forest products.

MAP 13:



Prior to this extensive investment, in 1997 the State of Vermont purchased easements on 31,450 acres in another major land deal with the former John Hancock Insurance properties creating Legacy Forests. While the easements limit future subdivision, they protect public

access and recreation opportunities. These new lands and previously state owned lands provide the Northeast Kingdom with some of the finest, most remote, and diverse outdoor recreation opportunities in New England.

Table 9.2: Public Lands in the NEK

| Town | Parcel Name | Acres |
|---------------------------------|------------------------------------|--------------|
| Averill | Averill Mountain WMA | 510 |
| Newark | Bald Hill Wildlife Management Area | 932 |
| Troy | Big Falls SP | 16 |
| Holland | Bill Sladyk WMA | 9,496 |
| Norton | Black Turn Brook SF | 593 |
| Brighton | Brighton SP | 152 |
| Sutton | Calendar Brook WMA | 340 |
| Barton | Crystal Lake SP | 16 |
| Burke | Darling State Park | 1,997 |
| Groton, Peacham | Groton SF | 23,706 |
| Burke | Hazens Notch SP | 307 |
| Sheffield | Holbrook SP | 202 |
| Jay | Jay SF | 3,877 |
| Peacham | Levi Pond WMA | 260 |
| Jay | Long Trail SF | 2,774 |
| Lyndon | Lyndon State Forest | 72 |
| Maidstone | Maidstone SF | 475 |
| Wheelock, Sheffield | Mathewson SF | 795 |
| Ryegate, Barnet | Roy Mountain WMA | 1,590 |
| Westmore | Sentinel Rock SP | 330 |
| Irasburg | South Bay WMA | 1,515 |
| Walden, Stannard, Wheelock | Steam Mill Brook | 10,421 |
| Victory | Victory Basin WMA | 4,970 |
| Victory, Lunenburg | Victory SF | 15,997 |
| Barton | Wenlock WMA | 1,994 |
| Brunswick, Ferdinand, Maidstone | West Mountain WMA | 22,738 |
| Barton | Willoughby Falls WMA | 130 |
| Westmore, Sutton | Willoughby SF | 7,300 |

(NVDA, 2002)

NATURAL RESOURCE GOALS

- The overarching goal for the region is to balance local economic needs with the protection of the natural resource that so many of the region's residents enjoy and depend upon.
- The quality and quantity of the region's surface waters should be protected, maintained, and restored.
- The quality and quantity of existing and potential groundwater resources should be protected and improved.
- Significant wetlands within the region should be protected.
- The region's mineral and soil resources should be used in a manner that will support the sustainable growth and development of the region.

- A consistently high level of air quality should be maintained for the health, safety, and enjoyment of the region's residents and visitors.
- Adequate resource information for the region should be maintained to improve the region's ability to plan for protection of wildlife resources in the area.
- Critical wildlife habitat should be protected.
- The native biodiversity of the region should be maintained, and restored when appropriate.
- Private, public and community interests should be considered in matters affecting local recreation and open space.

STRATEGIES

- Provide public education on state and local water quality issues as they relate to local planning and development.
- Discourage inappropriate development in flood hazard areas and floodplains. Support compatible land uses in flood areas, such as agriculture.
- Support the efforts of watershed organizations working in the region.
- Encourage and assist communities to identify and protect community water supplies. Education on water conservation and resource protection should accompany these efforts.
- Prevent the degradation of significant wetlands through public education.
- Minimize the negative impacts of mineral and earth resource extraction and processing facilities.
- Support development of new markets and uses for local mineral resources. Encourage the use of locally obtained minerals for building construction and highway construction and maintenance.
- Support efforts to reduce air pollutants generated in the region from the residential, commercial, industrial, and transportation sectors.
- Support broader state and regional efforts to minimize pollutants entering the region from out of state.
- Support local and state efforts that inventory, delineate and map important habitats and wetlands.
- Support local efforts to protect critical wildlife habitat and maintain habitat connectivity.
- Assist interested towns with planning and mapping for the protection of habitats and natural resources.
- Support state and local efforts to mitigate the impacts of the non-native species through ecologically sound methods (e.g. insect control, etc.).
- Support the protection of endangered and threatened native species.
- Maintain and improve the resource stewardship in the area by supporting and advocating for recreation and environmental education opportunities.
- Support the protection and the acquisition of unique and irreplaceable recreational spaces open for the public to enjoy.
- Encourage the growth of businesses focused on recreational activities. Support ecotourism in the region that will increase visits to the region and minimize the disturbance of important habitats.